# SANTA CRUZ BIOTECHNOLOGY, INC.

# CaMKII8 (L-04): sc-100362



## BACKGROUND

The Ca<sup>2+</sup>/calmodulin-dependent protein kinases (CaM kinases) comprise a structurally related subfamily of serine/threonine kinases which include CaMKI, CaMKII and CaMKIV. CaMKII is a ubiquitously expressed serine/ threonine protein kinase that is activated by Ca<sup>2+</sup> and calmodulin (CaM) and has been implicated in regulation of the cell cycle and transcription. There are four CaMKII isozymes, designated  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ , which may or may not be co-expressed in the same tissue type. CaMKIV is stimulated by Ca<sup>2+</sup> and CaM but also requires phosphorylation by a CaMK for full activation. Stimulation of the T cell receptor CD3 signaling complex with an anti-CD3 monoclonal antibody leads to a 10-40 fold increase in CaMKIV activity. An additional kinase, CaMKK, functions to activate CaMKI through the specific phosphorylation of the regulatory threonine residue at position 177.

## REFERENCES

- 1. Tombes, R.M., et al. 1995. G<sub>1</sub> cell cycle arrest apoptosis are induced in NIH 3T3 cells by KN-93, an inhibitor of CaMK-II (the multifunctional Ca<sup>2+</sup>/CaM kinase). Cell Growth Differ. 6: 1063-1070.
- 2. Baltas, L.G., et al. 1995. The cardiac sarcoplasmic reticulum phospholamban kinase is a distinct  $\delta$ -CaM kinase isozyme. FEBS Lett. 373: 71-75.

## **CHROMOSOMAL LOCATION**

Genetic locus: CAMK2D (human) mapping to 4q26; Camk2d (mouse) mapping to 3 G1.

#### SOURCE

 ${\rm CaMKII\delta}$  (L-04) is a mouse monoclonal antibody raised against recombinant CaMKII\delta of human origin.

#### PRODUCT

Each vial contains 100  $\mu g$   $lgG_{2b}$  lambda light chain in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

## **APPLICATIONS**

CaMKII $\delta$  (L-04) is recommended for detection of CaMKII $\delta$  of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CaMKII& siRNA (h): sc-38953, CaMKII& siRNA (m): sc-38954, CaMKII& siRNA (r): sc-270384, CaMKII& shRNA Plasmid (h): sc-38953-SH, CaMKII& shRNA Plasmid (m): sc-38954-SH, CaMKII& shRNA Plasmid (r): sc-270384-SHCaMKII& shRNA (h) Lentiviral Particles: sc-38953-V, CaMKII& shRNA (m) Lentiviral Particles: sc-38954-V and CaMKII& shRNA (r) Lentiviral Particles: sc-270384-V.

Molecular Weight of CaMKII8: 54 kDa.

Positive Controls: CaMKII $\delta$  (h): 293T Lysate: sc-115074 or mouse brain extract: sc-2253.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA





CaMKII& (L-04): sc-100362. Western blot analysis of CaMKII& expression in non-transfected: sc-117752 (A) and human CaMKII& transfected: sc-115074 (B) 293T whole cell lysates.

CaMKII8 (L-04): sc-100362. Immunofluorescence staining of paraformaldehyde-fixed NIH/3T3 cells showing cytoplasmic localization.

#### **SELECT PRODUCT CITATIONS**

- Ma, H., et al. 2014. γCaMKII shuttles Ca<sup>2+</sup>/CaM to the nucleus to trigger CREB phosphorylation and gene expression. Cell 159: 281-294.
- Wang, X., et al. 2018. Chemotherapy-induced differential cell cycle arrest in B-cell lymphomas affects their sensitivity to Wee1 inhibition. Haematologica 103: 466-476.
- 3. Nhieu, J., et al. 2020. Noncanonical retinoic acid signaling. Methods Enzymol. 637: 261-281.
- Dalal, P.J., et al. 2021. Spatiotemporal restriction of endothelial cell calcium signaling is required during leukocyte transmigration. J. Exp. Med. 218: e20192378.
- Zhang, X., et al. 2022. Interaction between A-kinase anchoring protein 5 and protein kinase A mediates CaMKII/HDAC signaling to inhibit cardiomyocyte hypertrophy after hypoxic reoxygenation. Cell. Signal. 103: 110569.
- Chuah, Y.H., et al. 2023. CAMK2D serves as a molecular scaffold for RNF8-MAD2 complex to induce mitotic checkpoint in glioma. Cell Death Differ. 30: 1973-1987.
- Ji, M., et al. 2023. CaMKII regulates the proteins TPM1 and MYOM2 and promotes diacetylmorphine-induced abnormal cardiac rhythms. Sci. Rep. 13: 5827.
- Chinnappa, S., et al. 2023. Beta blockade prevents cardiac morphological and molecular remodelling in experimental uremia. Int. J. Mol. Sci. 25: 373.
- Chai, B., et al. 2024. Tau aggregation-dependent lipid peroxide accumulation driven by the hsa\_circ\_0001546/14-3-3/CAMK2D/Tau complex inhibits epithelial ovarian cancer peritoneal metastasis. Adv. Sci. 11: e2310134.
- Manolis, D., et al. 2024. Quantitative proteomics reveals CLR interactome in primary human cells. J. Biol. Chem. 300: 107399.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.