

POPX2 (SJ-6): sc-100416

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine protein phosphatases. POPX1, also designated partner of PIX 1, PPM1E (protein phosphatase 1E) or PP2CH, and POPX2, also designated partner of PIX 2, PPM1F (protein phosphatase 1F), CaMKPase (CaM-kinase phosphatase), or FEM-2, belong to the PP2C family of serine/threonine phosphatases. Members of the PP2C family are negative regulators of cell stress response pathways. POPX2 is a ubiquitously expressed protein and POPX1 is predominantly expressed in brain and testis. POPX1 and POPX2 specifically interact with PIX (PAK interacting exchange factor) proteins and negatively regulate the activity of α PAK, a protein kinase that can lead to the breakdown of actin stress fibers and other morphological changes. POPX2 can also interact with and regulate CaMKII activity. Overexpression of POPX2 can result in caspase-dependent apoptosis.

REFERENCES

1. Nomura, N., et al. 1994. Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res.* 1: 27-35.
2. Kikuno, R., et al. 1999. Prediction of the coding sequences of unidentified human genes. XIV. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 6: 197-205.
3. Tan, K.M., et al. 2001. The *Caenorhabditis elegans* sex-determining protein FEM-2 and its human homologue, hFEM-2, are Ca^{2+} /calmodulin-dependent protein kinase phosphatases that promote apoptosis. *J. Biol. Chem.* 276: 44193-44202.
4. Koh, C.G., et al. 2002. The p21-activated kinase PAK is negatively regulated by POPX1 and POPX2, a pair of serine/threonine phosphatases of the PP2C family. *Curr. Biol.* 12: 317-321.
5. Harvey, B.P., et al. 2004. Regulation of the multifunctional Ca^{2+} /calmodulin-dependent protein kinase II by the PP2C phosphatase PPM1F in fibroblasts. *J. Biol. Chem.* 279: 24889-24898.
6. Ishida, A., et al. 2005. Identification of major Ca^{2+} /calmodulin-dependent protein kinase phosphatase-binding proteins in brain: biochemical analysis of the interaction. *Arch. Biochem. Biophys.* 435: 134-146.

CHROMOSOMAL LOCATION

Genetic locus: PPM1F (human) mapping to 22q11.22.

SOURCE

POPX2 (SJ-6) is a mouse monoclonal antibody raised against recombinant POPX2 of human origin.

PRODUCT

Each vial contains 100 μg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

POPX2 (SJ-6) is recommended for detection of POPX2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for POPX2 siRNA (h): sc-62844, POPX2 shRNA Plasmid (h): sc-62844-SH and POPX2 shRNA (h) Lentiviral Particles: sc-62844-V.


Molecular Weight of POPX2: 54 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204 or POPX2 (h2): 293T Lysate: sc-177761.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



POPX2 (SJ-6): sc-100416. Western blot analysis of POPX2 expression in non-transfected 293T: sc-117752 (A), human POPX2 transfected 293T: sc-177761 (B) and Jurkat (C) whole cell lysates.



POPX2 (SJ-6): sc-100416. Western blot analysis of POPX2 expression in non-transfected 293T: sc-117752 (A), human POPX2 transfected 293T: sc-177760 (B) and Jurkat (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.